

Math 107-04,07 Sample Final Exam, Spring 2009

Dr. Richard Mikula, Instructor

April 29, 2009

1 Descriptive Statistics

Given the data set

20, 50, 60, 70, 71, 72, 73, 73, 80, 89

Find:

1. The mean

Answer: 65.8

2. The median

Answer: 71.5

3. The standard deviation (by either technique) and the variance

Answers: 19.17058, 367.5111

4. The $100 \cdot p$ th percentile for

(a) $p = 0.1$

(b) $p = 0.17$

(c) $p = 0.25$

(d) $p = 0.5$

(e) $p = 0.75$

Answers: 35, 50, 60, 71.5, 73

5. Use a box and whisker argument to find any outliers.

Answer: 20 is an outlier, since it does not lie in the interval $[40.5, 92.5]$.

2 Probability

1. Suppose you toss a "fair" 20 sided die (in the shape of a icosahedron) with the faces labeled 1, 2, ... , 20. Find the probability that:

- (a) You roll a multiple of 3.
- (b) You don't roll a multiple of 3.

Answers: $\frac{6}{20}$ and $\frac{14}{20}$ respectively.

2. For the sample space

$$S = \{a, b, c, d\},$$

do the following tables define a probability distribution? Please explain.

outcome	probability
(a) a	0
b	0
c	0
d	1

outcome	probability
(b) a	-0.001
b	0
c	0.001
d	1

outcome	probability
(c) a	0.23
b	0.23
c	0.23
d	0.29

outcome	probability
(d) a	0.3
b	0.3
c	0.2
d	0.22

outcome	probability
(e) a	0.25
b	0.5
c	0.15
d	0.1

Answers: Yes, No, No, No and Yes respectively.

3. Suppose an experiment has two events A, B with the following probabilities

$$P(A) = \frac{1}{3}, P(B) = \frac{1}{2}, P(A \cap B^c) = \frac{1}{6}$$

Find:

- (a) $P(A \cap B)$
- (b) $P(A^c)$
- (c) $P(B^c)$
- (d) $P(A \cup B)$
- (e) $P(A^c \cup B^c)$

Answers: $\frac{1}{6}$, $\frac{2}{3}$, $\frac{1}{2}$, $\frac{2}{3}$, and $\frac{5}{6}$ respectively. To see these, use a Venn diagram as a visual aid.

4. Suppose A and B are independent events with probabilities:

$$P(A) = 0.3, P(B) = 0.6$$

Find:

- (a) $P(A \cap B)$
- (b) $P(A \cup B)$
- (c) $P(A^c \cup B^c)$
- (d) $P((A \cup B)^c)$
- (e) $P(A|B)$

Answers: 0.18, 0.72, 0.82, 0.28 and 0.3 respectively. To see these, use a Venn diagram as a visual aid.

5. Suppose A and B are events with

$$P(A) = 0.35, P(B) = 0.25, P(A \cap B) = 0.2$$

- (a) Find $P(A|B)$.
- (b) Find $P(B|A)$.
- (c) Are A, B independent events? Please explain.

Answers: 0.8, 0.5714, and No respectively.

6. In a math course suppose that 25 percent of the students failed the first exam, 15 percent failed the second exam and 5 percent failed both exams.

- (a) Are the events failing the first exam and failing the second exam independent?
- (b) Find $P(\text{Failed second exam given failed first exam})$.

Answers: No and 0.2 respectively.

3 Linear Regression

Given the data set

$$(1, 75), (2, 100), (3, 65), (4, 72), (6, 45)$$

Find:

1. The line of best-fit.
2. Pearson's correlation coefficient.

Answers: $S_{xx} = 14.8$ $S_{yy} = 1569.2$, $S_{xy} = -114.4$, $\bar{x} = 3.2$, $\bar{y} = 71.5$, $m = -7.72973$, $b = 96.73514$, $r = -0.75068$.

Thus $y = -7.72973x + 96.73514$.

4 Multivariate Categorical Data and Independence

A random group of 300 adult males was cross-classified according to age and cholesterol level. The following table has the results:

Age vrs Cholesterol level	Low	Medium	high cholesterol	total
20-34 yrs.	66	24	8	98
35-54 yrs.	54	48	22	124
55-74 yrs.	18	50	10	78
total	138	122	40	300

Using the χ^2 test for independence, at the $\alpha = 0.05$ level, test the hypothesis that age and cholesterol level are independent. **Answer:** $\chi^2 = 39.526$, and thus we reject H_0 in favor of H_1 : dependence.

5 Random Variables

1. Let X be a random variable with probability density function $f(x)$ given by

x	$f(x)$
0	0.1
1	0.1
2	0.4
3	0.15
4	0.15
5	0.1

Find:

- (a) $P(X \text{ is even })$
- (b) $E(X)$

(c) $sd(X)$

Answers: 0.65, 2.45, 1.40

2. Suppose 10 percent of the items produced by a factory are defective. Suppose that 8 are chosen at random. Let X be the number of defective items in the collection of 8 that were chosen. Find:

(a) $P(X = 4)$

(b) $P(X < 2)$

(c) $P(X \geq 2)$

(d) $P(X = 0)$

(e) $P(X > 0)$

(f) $E(X)$

(g) $sd(X)$

Answers: 0.0046, 0.1869, 0.4305, 0.5695, 0.8, 0.8485

3. If for the above factory, 100 items are chosen. Again letting X be the number of defective items, use Chebyshev's inequality to find a lower bound for

$$P(0 < X < 20).$$

Answer: 0.91

6 The Normal distribution

1. Suppose the heights of American males are (approximately) normally distributed with a mean $\mu = 68$ inches and standard deviation $\sigma = 2.5$ inches. Let $X(\text{male}) = \text{height}$ in inches. Find:

(a) $P(60 < X < 65)$

(b) $P(63 < X \leq 73)$

(c) $P(X \geq 72)$

(d) $P(X > 72)$

(e) $P(X \leq 65)$

Answers: 0.114, 0.954, 0.055, 0.055, 0.115